

Mass and angular momenta of Kerr anti-de Sitter spacetimes in Einstein-Gauss-Bonnet theory

Nathalie Deruelle* and Yoshiyuki Morisawa**

**Institut d'Astrophysique de Paris,
GReCO, FRE 2435 du CNRS,
98 bis boulevard Arago, 75014, Paris, France*
and

*Institut des Hautes Etudes Scientifiques,
35 Route de Chartres, 91440, Bures-sur-Yvette, France*

*** Yukawa Institute for Theoretical Physics
Kyoto University
Kyoto 606-8502, Japan*

29 November 2004

Abstract

We compute the mass and angular momenta of rotating anti-de Sitter spacetimes in Einstein-Gauss-Bonnet theory of gravity using a superpotential derived from standard Noether identities. The calculation relies on the fact that the Einstein and Einstein-Gauss-Bonnet vacuum equations are the same when linearized on maximally symmetric backgrounds and uses the recently discovered D-dimensional Kerr-anti-de Sitter solutions to Einstein's equations.

deruelle@ihes.fr
morisawa@yukawa.kyoto-u.ac.jp

In [1] Gibbons, Lu, Page and Pope found D-dimensional Kerr-anti-de Sitter solutions of Einstein's equations in vacuo. In [2] Gibbons, Perry and Pope calculated the angular momenta $J_{(D)i}$ of these rotating spacetimes using Komar's integrals. They also obtained the mass $E_{(D)}$ of rotating *black holes* using the first law of thermodynamics. In [3], the mass of these rotating spacetimes was computed using the covariant Katz-Bičák-Lynden Bell (KBL) superpotential [4], that is by means of classical Noether identities. The results confirm those of [2] and hold whether the source is a black hole or a "star" because they do not depend on the actual source of curvature.⁽¹⁾

In Einstein-Gauss-Bonnet (EGB) theory in 5 dimensions, the static spherically symmetric Schwarzschild-like solution was obtained in [5] and its mass was computed by various methods, including the use of the KBL superpotential extended to the Einstein-Gauss-Bonnet Lagrangian, see [6].

The rotating Kerr-like solution of the EGB equations is not known (apart from the topologically special solution given in [7]). However, on one hand, the Einstein and Einstein-Gauss-Bonnet vacuum equations are the same when linearized on a maximally symmetric background (see [5] and below) : hence the D-dimensional Kerr-anti-de Sitter solutions to the Einstein field equations found in [1] are also asymptotic solutions to the Einstein-Gauss-Bonnet equations. On the second hand, only the asymptotic form of the metric is required to define and compute the mass and angular momenta of a given spacetime when using superpotentials derived from Noether identities.

The object of this Note is to use these two remarks to compute the mass and angular momenta of Kerr-AdS spacetimes in EGB theory by means of the KBL superpotential proposed in [6].

The Einstein-Gauss-Bonnet Lagrangian can be written as (see e.g. [6])

$$L \equiv -2\Lambda + R + \alpha R^{\mu\nu\rho\sigma} P_{\mu\nu\rho\sigma} \quad (1)$$

with $P_{\mu\nu\rho\sigma} \equiv R_{\mu\nu\rho\sigma} - 2(R_{\mu[\rho}g_{\sigma]\nu} - R_{\nu[\rho}g_{\sigma]\mu}) + R g_{\mu[\rho}g_{\sigma]\nu}$

where Λ is a (negative) cosmological constant, α a coupling constant which is zero in Einstein's theory, and $R_{\mu\nu\rho\sigma}$, $R_{\mu\nu}$ and R the Riemann tensor, Ricci tensor and curvature scalar of the metric $g_{\mu\nu}$. Brackets stand for anti-symmetrization : $f_{[\mu\nu]} \equiv \frac{1}{2}(f_{\mu\nu} - f_{\nu\mu})$. The equations of motion derived from that Lagrangian are

$$R^\mu_\nu + 2\alpha R^{\mu\lambda\rho\sigma} P_{\nu\lambda\rho\sigma} - \frac{1}{2}\delta^\mu_\nu L = 0. \quad (2)$$

We linearize them, that is, we set

$$g_{\mu\nu} = \bar{g}_{\mu\nu} + h_{\mu\nu} + \mathcal{O}(h^2). \quad (3)$$

⁽¹⁾ In the second version of their paper, Gibbons Perry and Pope showed that the mass can also be obtained using the classical Ashtekar-Magnon-Das conformal boundary formula.

We choose the background $\bar{g}_{\mu\nu}$ to be the metric of a maximally symmetric spacetime $\bar{\mathcal{M}}$ solving the field equations (2). Since they are quadratic in the Riemann tensor there are two solutions, defined by

$$\bar{R}_{\mu\nu\rho\sigma} = -\frac{1}{\mathcal{L}^2}(\bar{g}_{\mu\rho}\bar{g}_{\nu\sigma} - \bar{g}_{\mu\sigma}\bar{g}_{\nu\rho}) \quad \text{where} \quad \frac{1}{\mathcal{L}^2} \equiv \frac{1}{2\tilde{\alpha}} \left(1 \pm \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \right) \quad (4)$$

with $\tilde{\alpha} \equiv (D-3)(D-4)\alpha$ and $l^2 \equiv -\frac{(D-1)(D-2)}{2\Lambda}$, D being the dimension of spacetime. We shall choose the lower sign for $\mathcal{L}^2$ so that the limit when $\alpha = 0$ solves the Einstein equations. (When Λ and α are such that $\sqrt{1 - 4\tilde{\alpha}/l^2} = 0$ the two roots coalesce into a single double root.) It is an exercise to show that at linear order the field equations (2) read (a result already obtained in [5] for $\Lambda = 0$)

$$\sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \left[R_{\nu}^{\mu} - \frac{1}{2}\delta_{\nu}^{\mu} R + \delta_{\nu}^{\mu} \Lambda \frac{l^2}{\mathcal{L}^2} \right] = \mathcal{O}(h^2) \quad (5)$$

where ($\bar{\nabla}_{\mu}$ being the covariant derivative associated to $\bar{g}_{\mu\nu}$)

$$R_{\nu}^{\mu} = -\frac{(D-1)}{\mathcal{L}^2}(\delta_{\nu}^{\mu} - h_{\nu}^{\mu}) + \frac{1}{2} \left[\bar{\nabla}_{\alpha}(\bar{\nabla}_{\nu} h^{\alpha\mu} + \bar{\nabla}^{\mu} h_{\nu}^{\alpha} - \bar{\nabla}^{\alpha} h_{\nu}^{\mu}) - \bar{\nabla}^{\mu} \bar{\nabla}_{\nu} h_{\rho}^{\rho} \right] + \mathcal{O}(h^2) \quad (6)$$

is the linearized Ricci tensor on $\bar{\mathcal{M}}$. Hence the linearized Einstein-Gauss-Bonnet equations are the same as the linearized pure Einstein equations with effective cosmological constant $\Lambda_e \equiv \Lambda \frac{l^2}{\mathcal{L}^2}$, and up to the overall factor $\sqrt{1 - 4\tilde{\alpha}/l^2}$ that we shall assume not to be zero.

The general D-dimensional Kerr-AdS metrics solutions to Einstein's equations found in [1] therefore also solve the EGB equations at linear order. As an example, the 5-dimensional Einstein-Kerr-AdS metric reads, in Kerr-Schild ellipsoidal coordinates $x^{\mu} = (t, r, \theta, \phi, \psi)$ [1] :

$$ds^2 = d\bar{s}^2 + \frac{2m}{U} (h_{\mu} dx^{\mu})^2 \quad (7)$$

with $d\bar{s}^2$ the AdS line element of $\bar{\mathcal{M}}$:

$$\begin{aligned} d\bar{s}^2 = & -\frac{(1 + r^2/\mathcal{L}^2)\Delta_{\theta}}{\Xi_a \Xi_b} dt^2 + \frac{r^2 \rho^2}{(1 + r^2/\mathcal{L}^2)(r^2 + a^2)(r^2 + b^2)} dr^2 + \frac{\rho^2}{\Delta_{\theta}} d\theta^2 + \\ & + \frac{r^2 + a^2}{\Xi_a} \sin^2 \theta d\phi^2 + \frac{r^2 + b^2}{\Xi_b} \cos^2 \theta d\psi^2 \end{aligned} \quad (8)$$

where $\Delta_{\theta} \equiv \Xi_a \cos^2 \theta + \Xi_b \sin^2 \theta$, where $\rho^2 \equiv r^2 + \mathcal{L}^2(1 - \Delta_{\theta})$, and where Ξ_a and Ξ_b are related to the rotation parameters a and b by $\Xi_a \equiv 1 - a^2/\mathcal{L}^2$, $\Xi_b \equiv 1 - b^2/\mathcal{L}^2$. As for the function U and the null vector h_{μ} they are given by $U = \rho^2$ and

$$h_{\mu} dx^{\mu} = \frac{\Delta_{\theta}}{\Xi_a \Xi_b} dt + \frac{r^2 \rho^2}{(1 + r^2/\mathcal{L}^2)(r^2 + a^2)(r^2 + b^2)} dr + \frac{a \sin^2 \theta}{\Xi_a} d\phi + \frac{b \cos^2 \theta}{\Xi_b} d\psi. \quad (9)$$

Note that the full Ricci tensor of the Kerr-Schild metric (7-9) is linear in $h_{\mu\nu} = \frac{2m}{U} h_\mu h_\nu$ and hence exactly given by (6) [1]. Note also [1] that the metric (7-9) tends for large r to the AdS metric in static (non rotating) coordinates.

There are various ways to associate conserved charges to the EGB Lagrangian (see e.g. [6] for a short review). We shall use here the superpotential $J^{[\mu\nu]}$ proposed in [6] which is defined as

$$\hat{J}^{[\mu\nu]} \equiv \hat{J}_E^{[\mu\nu]} + \alpha \hat{J}_{GB}^{[\mu\nu]} \quad (10)$$

where a hat means multiplication by $\sqrt{-g}$, with $g = \bar{g}$ the determinant of the metric (7-9). The Einstein contribution $\hat{J}_E^{[\mu\nu]}$ is given by, see [4]

$$\begin{aligned} -8\pi \hat{J}_E^{[\mu\nu]} &\equiv D^{[\mu} \hat{\xi}^{\nu]} - \overline{D^{[\mu} \hat{\xi}^{\nu]}} + \hat{\xi}^{[\mu} k_E^{\nu]} \\ \text{with } k_E^\nu &\equiv g^{\nu\rho} \Delta_{\rho\sigma}^\sigma - g^{\rho\sigma} \Delta_{\rho\sigma}^\nu \quad \text{and} \quad \Delta_{\nu\rho}^\mu \equiv \Gamma_{\nu\rho}^\mu - \overline{\Gamma_{\nu\rho}^\mu} \end{aligned} \quad (11)$$

where $\Gamma_{\nu\rho}^\mu$ are the Christoffel symbols of metric (7-9) and where an overline means that the quantity is evaluated on the AdS spacetime $\overline{\mathcal{M}}$ (that is, by setting $m = 0$ in (7)). The following proposal for the Gauss-Bonnet contribution $\hat{J}_{GB}^{[\mu\nu]}$ was put forward in [6] :

$$\begin{aligned} -8\pi \hat{J}_{GB}^{[\mu\nu]} &\equiv 2 \left[P^{\mu\nu\alpha\beta} D_{[\alpha} \hat{\xi}_{\beta]} - \overline{P^{\mu\nu\alpha\beta} D_{[\alpha} \hat{\xi}_{\beta]}} \right] + \hat{\xi}^{[\mu} k_{GB}^{\nu]} \\ \text{with } k_{GB}^\nu &\equiv 4 \left(P^{\nu\alpha\beta}{}_\gamma - Q^{\nu\alpha\beta}{}_\gamma \right) \Delta_{\alpha\beta}^\gamma \end{aligned} \quad (12)$$

where the tensor $P_{\mu\nu\rho\sigma}$ is defined in (1) and where $Q_{\mu\nu\rho\sigma} \equiv 2(R_{\rho[\mu} g_{\nu]\sigma} + R_{\sigma[\mu} g_{\nu]\rho})$. When the vector ξ^μ is the Killing vector associated with time translations with respect to a non rotating frame at infinity, that is $\xi^\mu = (1, 0, 0, 0, 0)$, the associated conserved charge is the mass

$$M = \int_S d^{D-2} x \hat{J}^{[01]} \quad (13)$$

where S is the $(D-2)$ -sphere at infinity. If the vector ξ^μ is the Killing vector associated with rotations with respect to the ϕ (resp. ψ) axis, that is $\xi_a^\mu = (0, 0, 0, 1, 0)$ (resp. $\xi_b^\mu = (0, 0, 0, 0, 1)$), then the associated charges are the angular momenta.

To obtain the mass and angular momenta of the 5D-Kerr-like solution of EGB theory we inserted the metric (7-9) in the definitions (10-13) and took the large r limit (using *Maple* and *GR tensor*). The result is

$$M_{(5)} = \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \left[\frac{m\pi}{4(\Xi_a \Xi_b)^2} (2\Xi_a + 2\Xi_b - \Xi_a \Xi_b) \right] , \quad J_{(5)a} = \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \left[\frac{\pi m a}{2\Xi_a^2 \Xi_b} \right] . \quad (14)$$

We calculated similarly the mass and angular momentum in $D = 6, 7, 8$ dimensions in the case when there is only one non zero rotation parameter. The result is

$$M_D = \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \frac{m \mathcal{V}_{D-2}}{4\pi \Xi_a^2} \left[1 + \frac{(D-4)}{2} \Xi_a \right] , \quad J_D = \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \frac{m a \mathcal{V}_{D-2}}{4\pi \Xi_a^2} \quad (15)$$

where $\mathcal{V}_{D-2}$ is the volume of the $(D-2)$ -unit sphere. Finally, if one extrapolates to all dimensions the asymptotic expressions for the Kerr-AdS Riemann tensor obtained with *Maple* and *GR tensor* in $D = 5, 6, 7, 8$ dimensions,⁽²⁾ one finds (by hand) the mass and angular momenta of the general D-dimensional Kerr-like solutions of Einstein-Gauss-Bonnet theory as

$$M_{2n} = \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \left[\frac{m\mathcal{V}_{2n-2}}{4\pi\Xi} \sum_{i=1}^{i=n-1} \frac{1}{\Xi_i} \right], \quad M_{2n+1} = \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \left[\frac{m\mathcal{V}_{2n-1}}{4\pi\Xi} \left(\sum_{i=1}^{i=n} \frac{1}{\Xi_i} - \frac{1}{2} \right) \right]$$

$$J_{(D)i} = \sqrt{1 - \frac{4\tilde{\alpha}}{l^2}} \left[\frac{m\mathcal{V}_{D-2}}{4\pi\Xi} \frac{a_i}{\Xi_i} \right]. \quad (16)$$

Note that neither the background nor the vectors k_E^μ and k_{GB}^μ enter the calculation of the angular momenta. However they are not given simply by Komar's integrals because the first term in (12) is not zero. Expressions (14-16) have the right limits. When $\alpha = 0$ (Einstein's theory) they reduce to the results obtained in [2] and [3]. When there is no rotation ($a_i = 0, \Xi_i = 1$) the masses (16) reduce to the result obtained in [8] and [6] (noting that the coefficients of the metric (7) then tend to $g_{tt} \simeq 1/g_{rr} \simeq r^2/\mathcal{L}^2 - 2m/r^{D-3}$).⁽³⁾

The decisive check of formulae (14-16), that is of the pertinence of the proposal (12) made in [6] for the vector k_{GB}^μ , will be possible when the full geometry of the Kerr-like rotating *black holes* is known, by seeing if, with such definitions, the first law of thermodynamics is still satisfied.

Acknowledgments

N.D. thanks Joseph Katz for discussions and the Yukawa Institute for Theoretical Physics in Kyoto for hospitality and financial support. Y.M. is supported by a Grant-in-Aid for the 21st Century COE "Center for Diversity and Universality in Physics".

References

- [1] G.W. Gibbons, H. Lu, Don N. Page and C.N. Pope, *The General Kerr-de Sitter Metrics in All Dimensions*, hep-th/0404008.
- [2] G.W. Gibbons, M.J. Perry and C.N. Pope, *The First Law of Thermodynamics for Kerr-de Sitter Metrics*, hep-th/0408217.

⁽²⁾ for example : $\lim_{r \rightarrow \infty} (R_{101}^0 - \overline{R_{101}^0}) = \frac{m\mathcal{L}^2}{r^{D+1}}(D-1)[(D-3)W - 1]$, where $W \equiv \sum_{i=1}^{i=[D/2]} \frac{\mu_i^2}{\Xi_i}$ with $\sum_{i=1}^{i=[D/2]} \mu_i^2 = 1$ and $\Xi_{D/2} = 1$ if D is even.

⁽³⁾ Since the overall factor in (14-16) is the same as the one which appears in (5) one may conjecture that in the general Lovelock theory in D dimensions (whose Lagrangian is the sum of the first $[D/2]$ dimensionally continued Euler forms) the expressions for the masses and angular momenta will also be proportional to their values in Einstein's theory with an overall coefficient which is zero when the equation for the maximally symmetric space curvature has one single root of maximal multiplicity $[D/2]$.

- [3] N. Deruelle and J.Katz, *On the mass of a Kerr-anti-de Sitter spacetime in D dimensions*, gr-qc/0410135
- [4] J. Katz, *A note on Komar's anomalous factor*, Class. Quantum Grav., **2** (1985) 423; J. Katz, J. Bičák, D. Lynden-Bell, *Relativistic conservation laws and integral constraints for large cosmological perturbations*, Phys. Rev. **D55** (1997) 5759.
- [5] D.G. Boulware and S. Deser, *String generated gravity models*, Phys. Rev. Lett. **55** (1985) 2656
- [6] N. Deruelle, J. Katz and S. Ogushi, *Conserved Charges in Einstein- Gauss-Bonnet theory*, Class. Quantum Grav., **21** (2004) 1971, gr-qc/0310098.
- [7] M.H. Dehghani, *Charged Rotating Black Branes in anti-de Sitter Einstein-Gauss-Bonnet Gravity*, Phys.Rev. **D67** (2003) 064017, hep-th/0211191
- [8] Rong-Gen Cai, *Gauss-Bonnet Black Holes in AdS Spaces*, Phys.Rev. **D65** (2002) 084014, hep-th/0109133; I. Neupane, *Black hole entropy in string-generated gravity models*, Phys.Rev.**D67** (2003) 061501, hep-th/0212092